

Article

Deciphering the Molecular Mechanism Responsible for Efficiently Inhibiting Metastasis of Human Non-Small Cell Lung and Colorectal Cancer Cells Targeting the Matrix Metalloproteinases by *Selaginella repanda*

Adnan, Mohd, Siddiqui, Arif Jamal, Hamadou, Walid Sabri, Snoussi, Mejd, Badraoui, Riadh, Ashraf, Syed Amir, Jamal, Arshad, Awadelkareem, Amir Mahgoub, Sachidanandan, Manojkumar, Hadi, Ss, Khan, Mushtaq Ahmad and Patel, Mitesh

Available at <http://clock.uclan.ac.uk/37827/>

Adnan, Mohd, Siddiqui, Arif Jamal, Hamadou, Walid Sabri, Snoussi, Mejd, Badraoui, Riadh, Ashraf, Syed Amir, Jamal, Arshad, Awadelkareem, Amir Mahgoub, Sachidanandan, Manojkumar et al (2021) Deciphering the Molecular Mechanism Responsible for Efficiently Inhibiting Metastasis of Human Non-Small Cell Lung and Colorectal Cancer Cells Targeting the Matrix Metalloproteinases by Selaginella repanda. Plants, 10 (5). e979.

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.3390/plants10050979>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the policies page.



University of
Central Lancashire
UCLan

CLoK

Central Lancashire online Knowledge

www.clok.uclan.ac.uk



Article

Deciphering the Molecular Mechanism Responsible for Efficiently Inhibiting Metastasis of Human Non-Small Cell Lung and Colorectal Cancer Cells Targeting the Matrix Metalloproteinases by *Selaginella repanda*

Mohd Adnan ¹, Arif Jamal Siddiqui ¹, Walid Sabri Hamadou ¹, Mejdí Snoussi ¹, Riadh Badraoui ^{1,2}, Syed Amir Ashraf ³, Arshad Jamal ¹, Amir Mahgoub Awadelkareem ³, Manojkumar Sachidanandan ⁴, Sibte Hadi ⁵, Mushtaq Ahmad Khan ^{6,*} and Mitesh Patel ^{7,*}

- ¹ Department of Biology, College of Science, University of Hail, Hail P.O. Box 2440, Saudi Arabia; drmohtadnan@gmail.com (M.A.); arifjamal13@gmail.com (A.J.S.); walidsabrimail@gmail.com (W.S.H.); snmejd@yahoo.fr (M.S.); badraoui@yahoo.fr (R.B.); arshadjamalus@yahoo.com (A.J.)
- ² Section of Histology-Cytology, Medicine Faculty of Tunis, University of Tunis El Manar, La Rabta-Tunis 1007, Tunisia
- ³ Department of Clinical Nutrition, College of Applied Medical Sciences, University of Hail, Hail P.O. Box 2440, Saudi Arabia; amirashrafy2007@gmail.com (S.A.A.); mahgoubamir22@gmail.com (A.M.A.)
- ⁴ Department of Oral Radiology, College of Dentistry, University of Hail, Hail P.O. Box 2440, Saudi Arabia; smanojk68@gmail.com
- ⁵ School of Forensic and Applied Sciences, University of Central Lancashire, Preston PR1 2HE, UK; shadi@uclan.ac.uk
- ⁶ Department of Microbiology and Immunology, College of Medicine and Health Sciences, UAE University, Al Ain 17666, United Arab Emirates
- ⁷ Bapalal Vaidya Botanical Research Centre, Department of Biosciences, Veer Narmad South Gujarat University, Surat 394230, India
- * Correspondence: mushtaq.khan@uaeu.ac.ae (M.A.K.); patelmeet15@gmail.com (M.P.)



Citation: Adnan, M.; Siddiqui, A.J.; Hamadou, W.S.; Snoussi, M.; Badraoui, R.; Ashraf, S.A.; Jamal, A.; Awadelkareem, A.M.; Sachidanandan, M.; Hadi, S.; et al. Deciphering the Molecular Mechanism Responsible for Efficiently Inhibiting Metastasis of Human Non-Small Cell Lung and Colorectal Cancer Cells Targeting the Matrix Metalloproteinases by *Selaginella repanda*. *Plants* **2021**, *10*, 979. <https://doi.org/10.3390/plants10050979>

Academic Editor: Filippo Maggi

Received: 28 March 2021

Accepted: 8 May 2021

Published: 14 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: *Selaginella* species are known to have antimicrobial, antioxidant, anti-inflammatory, anti-diabetic as well as anticancer effects. However, no study has examined the cytotoxic and anti-metastatic efficacy of *Selaginella repanda* (*S. repanda*) to date. Therefore, this study aimed to evaluate the potential anti-metastatic properties of ethanol crude extract of *S. repanda* in human non-small-cell lung (A-549) and colorectal cancer (HCT-116) cells with possible mechanisms. Effect of *S. repanda* crude extract on the growth, adhesion, migration and invasion of the A-549 and HCT-116 were investigated. We demonstrated that *S. repanda* crude extract inhibited cell growth of metastatic cells in a dose and time dependent manner. Incubation of A-549 and HCT-116 cells with 100–500 µg/mL of *S. repanda* crude extract significantly inhibited cell adhesion to gelatin coated surface. In the migration and invasion assay, *S. repanda* crude extract also significantly inhibited cellular migration and invasion in both A-549 and HCT-116 cells. Moreover, reverse transcription-polymerase chain reaction, and real-time PCR (RT-PCR) analysis revealed that the activity and mRNA level of matrix metalloproteinase-9 (MMP-9), matrix metalloproteinase-2 (MMP-2) and membrane type 1-matrix metalloproteinase (MT1-MMP) were inhibited. While the activity of tissue inhibitor matrix metalloproteinase 1 (TIMP-1); an inhibitor of MMPs was stimulated by *S. repanda* crude extract in a concentration-dependent manner. Therefore, the present study not only indicated the inhibition of motility and invasion of malignant cells by *S. repanda*, but also revealed that such effects were likely associated with the decrease in MMP-2/-9 expression of both A-549 and HCT-116 cells. This further suggests that *S. repanda* could be used as a potential source of anti-metastasis agent in pharmaceutical development for cancer therapy.

Keywords: *Selaginella repanda*; non-small cell lung cancer; colorectal cancer; matrix metalloproteins; tissue inhibitor matrix metalloproteinase; metastasis; gene expression

1. Introduction

Over the past few years, cancer is the leading cause of death globally, resulting in about 9.6 million deaths in 2018. In men, commonly occurring cancers are lung, prostate, colorectal, stomach and liver, whereas, in women, cervical, lung, colorectal and thyroid cancers are most common [1]. Metastasis is the most devastating hallmark of cancer with more than 90% of deaths worldwide and a major obstacle in the treatment of cancers, which is defined as the emergence of secondary and tertiary tumors in tissues and organs aside from the origin [2,3]. Options available for the treatment of various types of cancers are chemotherapy, radiation therapy and surgery. All of these treatment options have harmful and severe adverse effects [4,5]. Therefore, in recent times, back to nature approach is highly accepted. A large number of plant-based molecules are developed and approved for anticancer and anti-metastatic therapies [6,7]. Hence, to find out the feasible course of action to have safe and to lower the adverse effects prompted by chemotherapy, the continuous exploration for anticancer phytochemicals plays a critical role.

The cosmopolitan genus *Selaginella* P. Beauv. also acknowledged as a “spike moss” belongs to the family *Selaginellaceae*, possessing about 700 to 750 species distributed around the globe. The *Selaginella* plants are usually used by a tribal community to cure sore throat, hepatic disorders, fever, cirrhosis, jaundice, cholecystitis, cough of lungs, diarrhea and various other related ailments. Moreover, it is also used for promoting the blood circulation, removing blood stasis, prevention of external bleeding after separation of the umbilical cord and trauma [8,9]. This is due to the presence of high content of different phytochemicals including, carbohydrates, chromones, benzenoids, alkaloids, coumarins, lignans, flavonoids, phenylpropanoids, steroids, pigments and quinoids [10–19]. Although the members of *Selaginella* are reported for their efficient medicinal applications, the cytotoxicity and anti-metastatic activity of *S. repanda* (Desv. & Poir.) against lung and colon cancers have not been described. Furthermore, matrix metalloproteinases (MMPs) are believed to be intricately involved in cancer development and progression. MMPs are known to play a vital role in tumor metastasis, angiogenesis and invasion, and are considered as a potential anticancer target. Therefore, the present study aimed to find out the cytotoxic and anti-metastatic efficacy of traditionally treasured plant *S. repanda* on human non-small cell lung and colorectal cancer with possible mechanisms explaining the expression of MMP-2/-9, MT1-MMP and TIMP-1 with and without treatment with *S. repanda*.

2. Results

2.1. High Resolution-Liquid Chromatography-Mass Spectroscopy (HR-LC-MS) Identification of *S. repanda* Phytoconstituents

Comprehensive phytoconstituent analysis from crude extract of *S. repanda* was carried out to analyze the plant's total extract obtained from 85% aqueous ethanol obtained from the whole plant of *S. repanda* via UHPLC-PDA-ESI-MS/MS. A diverse class of bioactive compounds was revealed, which are known to be potent and linked to various biological activities. Table 1 listed the identified phytoconstituents in *S. repanda* crude extract and are in agreement with previous studies [20,21]. A total of fifty-four constituents were tentatively identified. The molecular mass, mass spectral fragmentation patterns of the constituents, and standard samples were used for the purpose. Out of all the identified compounds, phenols and flavonoids were the major constituents. Identified phenols were isoferulic acid, chlorogenic acid, 4-coumaric acid, 7-hydroxycoumarine, coumarin, caffeic acid, formononetin, 4-methoxycinnamic acid and 2-amino-1,3,4-octadecanetriol. Identified potent flavonoid compounds were glycitin, vitexin, rhamnetin, kaempferol, luteolin, rutin, diosmetin, apigenin H, genistein. Identified potent terpenoids, alkaloids, benzenoids, monocarboxylic acid and ester compounds were urocanic acid, guvacoline, hordenine, sedanolide, scopoletin, norharman, citral, methylcinnamate, 8-hydroxyquinoline, pulegone and 4-hydroxyphenylacetic acid [9].

Table 1. Phytochemical composition of *S. repanda* crude extract identified by HR-LC-MS technique [9].

Phytocompounds	Formula	Class	<i>m/z</i>	RT (min)	Mass
Urocanic acid	C ₆ H ₆ N ₂ O ₂	monocarboxylic acid	131.2	24.845	138.04267
Hordeanine	C ₁₀ H ₁₅ NO	alkaloid	167.2	23.071	169.23958
Oleamide	C ₁₈ H ₃₅ NO	fatty acid	286.1	20.177	281.27135
Hexadecanamide	C ₁₆ H ₃₃ NO	fatty acid amide	250.4	19.849	255.25575
Arachidonic acid	C ₂₀ H ₃₂ O ₂	polyunsaturated fatty acid	301.8	18.601	304.2397
4-Coumaric acid	C ₉ H ₈ O ₃	phenol	168.2	18.265	164.04718
2-Arachidonoyl glycerol	C ₂₃ H ₃₈ O ₄	fatty acid derivative	371.0	16.012	378.27635
Valine	C ₅ H ₁₁ NO ₂	amino acid	118.4	13.901	117.07901
Genistein	C ₁₅ H ₁₀ O ₅	isoflavone	275.3	13.542	270.05214
Diosmetin	C ₁₆ H ₁₂ O ₆	flavonoid	294.3	13.534	300.06279
2-Amino-1,3,4-octadecanetriol	C ₁₈ H ₃₉ NO ₃	phenol	311.5	13.499	317.29231
Glycitein	C ₁₆ H ₁₂ O ₅	flavonoid	288.9	13.477	284.06807
Rhamnetin	C ₁₆ H ₁₂ O ₇	flavonoid	311.2	12.452	316.05766
Formononetin	C ₁₆ H ₁₂ O ₄	phenol	265.3	12.157	268.07329
Luteolin	C ₁₅ H ₁₀ O ₆	flavonoid	278.6	11.735	286.04723
Apigenin	C ₁₅ H ₁₀ O ₅	flavone	272.4	11.177	270.05235
Glycitin	C ₂₂ H ₂₂ O ₁₀	isoflavone	442.2	11.088	446.12065
(-)-Caryophyllene oxide	C ₁₅ H ₂₄ O	epoxide	228.4	10.861	220.18227
Kuromanin	C ₂₁ H ₂₀ O ₁₁	pigment	458.1	8.976	448.09996
Kaempferol	C ₁₅ H ₁₀ O ₆	flavonoid	294.6	8.662	286.04726
Sedanolid	C ₁₂ H ₁₈ O ₂	isobenzofuran	199.5	8.578	194.13039
α-Pinene-2-oxide	C ₁₀ H ₁₆ O	terpenoid	148.9	8.494	152.11989
Quercetin-3β-D-glucoside	C ₂₁ H ₂₀ O ₁₂	flavonoid	260.3	8.438	464.09465
Quercetin	C ₁₅ H ₁₀ O ₇	flavonoid	308.6	8.414	302.04192
Vitexin	C ₂₁ H ₂₀ O ₁₀	flavonoid	436.8	8.319	432.10497
Rutin	C ₂₇ H ₃₀ O ₁₆	flavonoid	615.4	8.291	610.15239
4-Methoxycinnamic acid	C ₁₀ H ₁₀ O ₃	phenol	186.2	7.754	178.06275
Norharman	C ₁₁ H ₈ N ₂	alkaloid	164.7	6.725	168.06847
4-Hydroxycoumarin	C ₉ H ₆ O ₃	benzopyrone	158.8	6.568	162.0314
Methyl cinnamate	C ₁₀ H ₁₀ O ₂	cinnamic acid ester	164.5	6.458	162.06775
Isoferulic acid	C ₁₀ H ₁₀ O ₄	phenol	198.6	6.367	194.05762
Scopoletin	C ₁₀ H ₈ O ₄	coumarin	196.3	5.969	192.04198
Citral	C ₁₀ H ₁₆ O	terpenoid	145.8	5.575	152.11989
Pulegone	C ₁₀ H ₁₆ O	terpenoid	158.9	4.94	152.11989
Caffeic acid	C ₉ H ₈ O ₄	phenol	186.2	4.692	180.04181
7-Hydroxycoumarin	C ₉ H ₆ O ₃	phenol	169.4	4.651	162.0314
Chlorogenic acid	C ₁₆ H ₁₈ O ₉	phenol	360.5	4.646	354.09435
Kynurenic acid	C ₁₀ H ₇ NO ₃	quinoline carboxylic acid	181.7	3.823	189.04239
Coumarin	C ₉ H ₆ O ₂	phenol	148.6	3.784	146.0365
3-Methylcrotonylglycine	C ₇ H ₁₁ NO ₃	amino acid	152.6	3.325	157.0737

Table 1. Cont.

Phytocompounds	Formula	Class	m/z	RT (min)	Mass
4-Hydroxyphenylacetic acid	C ₈ H ₈ O ₃	benzenoid	156.3	3.314	152.04714
8-Hydroxyquinoline	C ₉ H ₇ NO	alkaloid	153.2	2.965	145.05255
Maltol	C ₆ H ₆ O ₃	sugar	130.4	2.278	126.03161
Guvacoline	C ₇ H ₁₁ NO ₂	pyridine alkaloid	145.2	1.539	141.07878
L-Phenylalanine	C ₉ H ₁₁ NO ₂	amino acid	164.2	1.375	165.07883
L-Norleucine	C ₆ H ₁₃ NO ₂	amino acid	136.5	1.134	131.09453
L-Pyroglutamic acid	C ₅ H ₇ NO ₃	amino acid	133.1	1.04	129.0425
D-Glucosamine	C ₆ H ₁₃ NO ₅	amino sugar	183.3	0.946	179.079
Betaine	C ₅ H ₁₁ NO ₂	amino acid	122.2	0.935	117.07901
L(-)-Carnitine	C ₇ H ₁₅ NO ₃	amino acid derivative	157.6	0.93	161.10489
Acetylcholine	C ₇ H ₁₅ NO ₂	essential nutrient (vitamin)	149.7	0.850	145.11
α-Lactose	C ₁₂ H ₂₂ O ₁₁	sugar	349.9	0.839	342.11521
Choline	C ₅ H ₁₃ NO	essential nutrient (vitamin)	111.4	0.798	103.09988

Overall, LC-MS analysis endorsed the rich nature of the *S. repanda* plant in terms of varying chemical classes of compounds present in it, whereby the majority of all the identified compounds are flavonoids and phenolics in nature as previously described [9,20,21]. Flavonoids and natural phenolic acids are one of the most prevalent and pharmacologically active groups of plant secondary metabolites. Whereas, alkaloids are nitrogenous compounds that are widely distributed from prokaryotes to eukaryotes and are well-known for their different biological activities like anti-microbial, anti-HIV, anticancer and antiparasitic. These compounds play a crucial role in the prevention of cancer via various mechanisms at a molecular level. This may include impeding the signaling pathways, migration, differentiation and proliferation inhibition, gene regulation, carcinogen metabolism, induction of apoptosis via arresting cell cycle, etc. [22,23]. Therefore, the presence of diverse biologically active constituents contributed immensely to the plant's bioactive potential, and perhaps the ensuing better anticancer and anti-metastatic bioactivities.

2.2. Cytotoxic Effect of *S. repanda* Crude Extract

Cytotoxicity of *S. repanda* crude extract was evaluated against A549 and HCT-116 cancer cell lines by MTT assay at 24 and 48 h. Significant inhibition of both cancer cells' viability was observed in a time and dose-dependent manner. The IC₅₀ values for 24 h were found to be 341.1 µg/mL and 378.8 µg/mL (Figure 1C) and for 48 h were found to be 326.7 µg/mL and 351.4 µg/mL for A549 and HCT-116 cells (Figure 1D).

2.3. Anti-Migratory Effect of *S. repanda* Crude Extract

The most important metastatic event is the motility of cells, which takes place in different epithelial cells during the progression of cancer. Wound closure assay was carried out to study the effect of *S. repanda* crude extract on the migration of cells over wound scratch, made on culture plates. The migration of both A549 and HCT-116 cancer cells was evidently inhibited by the treatment of *S. repanda* crude extract in a time as well as dose-dependent manner, when compared to untreated control cells. Hence, *S. repanda* crude extract evidently inhibited the anchorage and spread of both cell lines along the edge of wound scratch (Figure 2A,B).




Figure 1. *S. repanda* plant and its anticancer activity. (A) Plant in wild (B) Close view of plant (C) Anticancer activity of *S. repanda* crude extract against A-549 and HCT-116 cancer cells for 24 h. (D) Anticancer activity of *S. repanda* crude extract against A-549 and HCT-116 cancer cells for 48 h. Error bars indicate SDs (\pm standard deviation) of three independent experiments. Significance; ns > 0.05, * p < 0.05, ** p < 0.005, *** p < 0.0005.

2.4. Anti-Invasion Effect of *S. repanda* Crude Extract

Transwell® cell culture inserts coated with Matrigel matrix were used to evaluate the ability of *S. repanda* crude extract to inhibit the invasion of A549 and HCT-116 cells, following treatment with different concentrations (100–500 µg/mL). The cellular invasion of both A549 and HCT-116 cancer cells was significantly inhibited by the treatment of *S. repanda* crude extract (100 µg/mL) in a time as well as dose-dependent manner, when compared to untreated control cells (Figure 2 C).

2.5. Anti-Adhesion Effect of *S. repanda* Crude Extract

Adhesion to endothelial cells by malignant cells is known to be mediated by extracellular matrix proteins (ECM), thereby, we evaluated the adhesion property of both A549 and HCT-116 cancer cells to gelatin coated surfaces both in the presence and absence of *S. repanda* crude extract. The adhesion of A549 and HCT-116 cells were reduced by 75.67% and 72.0% (500 µg/mL) at the incubation of 6 h. Relative adherence was measured by setting the number of adherent cells at 6 h to 100% (Figure 3A,B).




Figure 2. Anti-migration and anti-invasion effects of *S. repanda* crude extract. (A) Anti-migration activity on A-549 cancer cells (B) Anti-migration activity on HCT-116 cancer cells. The number of migrated cells was quantified in five different fields from three independent experiments. (C) Anti-invasion activity on A-549 and HCT-116 cancer cells. The invading cells were counted in five random fields under microscopes. Error bars indicate SDs (\pm standard deviation) of three independent experiments. Significance; ns > 0.05, * p < 0.05, ** p < 0.005, *** p < 0.0005.




Figure 3. Anti-adhesion effects of *S. repanda* crude extract on gelatin coated surfaces. (A) Anti-adhesion activity on A-549 cancer cells. (B) Anti-adhesion activity on HCT-116 cancer cells. Error bars indicate SDs (\pm standard deviation) of three independent experiments. Significance; ns > 0.05 , * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.

2.6. Changes in Transcriptional Level of Metastasis Related Genes

The expression level of metastasis related genes MMP-2, MMP-9, MT1-MMP and TIMP-1 in both A549 and HCT-116 malignant cells, which were induced by *S. repanda* crude extract were determined by real time PCR. Firstly, the expression level of MMP-2 and MMP-9 genes were evaluated and was found to be decreased in both cells treated with *S. repanda* crude extract after 24 h incubation, when compared to untreated cells (Figure 4A–D). After the analysis of MMP-2 and MMP-9, we further examined the effect of *S. repanda* crude extract on the expression of activator and inhibitor genes of MMPs (MT1-MMP and TIMP-1). The expression level of TIMP-1 was found to be increased and of MT1-MMP gene was decreased in both cells after the treatment of *S. repanda* crude extract, when compared to untreated cells after 24 h incubation (Figure 5A,B). These results show that *S. repanda* crude extract can regulate the expression of MMP-2 and MMP-9 genes, which can possibly further control the cascade of metastasis.




Figure 4. Effect of *S. repanda* crude extract on metastasis related genes. (A) Reverse transcription PCR to analyze the mRNA expression level of MMP-2 and MMP-9 in A-549 cancer cells. (B) Quantitative real-time PCR. (C) Reverse transcription PCR to analyze the mRNA expression level of MMP-2 and MMP-9 in HCT-116 cancer cells. (D) Quantitative real-time PCR. Significance; ns > 0.05, * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.




Figure 5. Effect of *S. repanda* crude extract on metastasis related genes. (A) Reverse transcription PCR to analyze the mRNA expression level of MT1-MMP and TIMP-1 in A-549 cancer cells. (B) Quantitative real-time PCR. (C) Reverse transcription PCR to analyze the mRNA expression level of MT1-MMP and TIMP-1 in HCT-116 cancer cells. (D) Quantitative real-time PCR. Significance; ns > 0.05, * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$.

3. Discussion

Regardless of all current progress in oncology, cancer is still one of the most life-threatening diseases around the globe [1]. It takes place as a localized disease, but can extend to different sites of the human body through migration, invasion and metastasis [2]. Metastasis is a multifaceted process involving an array of complicated mechanisms, which begins with extrication, accumulation and motility of cancer cells, followed by sticking to endothelial cells and the start of cancerous growth at different sites [24]. Progression of metastasis occurs after the degradation of ECM with cancer cells via different proteases like serine proteinase, cathepsins, MMPs and plasminogen activator, which prompts the separation of the intercellular matrix to promote the mobility of cancer cells [25]. Amongst involved proteases, MMP-9 and MMP-2 are profoundly implicated in cancer invasion and metastasis as they are the most essential for the degradation of base membranes [26–28].

Due to resistance to apoptosis and cytotoxic agents, metastasis is the dominant basis for cancer associated death. The percentage of morbidity and mortality in metastatic cancer patients is high, due to the failure of current chemotherapy agents to selectively and effectively kill cancer cells without destroying healthy cells at the sites of metastasis [29]. Metastasis is still a crucial clinical challenge in cancer treatment for researchers around the globe. Up till now, no such strong cleanse for cancer and its other catastrophic presentations have been found [29]. Currently, radiation therapy, chemotherapy, surgery are the conventional treatments for cancer. These treatments come with a range of side effects to human health; therefore, the importance of traditional medicines may decline. Thus, metastasis found the greatest challenging obstacle for successful cancer management and can be viewed as the last edge of cancer research [24].

Since ancient times, natural products have been recognized as an excellent source of bioactive compounds. They have been the primary source of general medicines, and can also be used directly as medicines [30–35]. Therefore, the pursuit and curiosity in the identification of medicinal plants and their derived natural products for evolving the novel cancer therapeutic strategies expanded vastly in recent times. In our previous study [9], a comprehensive phytochemical analysis from the crude extract of *S. repanda* was carried out via HR-LC-MS. A chromatogram was obtained with both positive and negative run and different types of phytochemicals were identified. In-depth profiling revealed various classes of metabolites such as alkaloids, flavonoids, sugars, vitamins, amino acids, phenols, terpenoids, phenols, etc. Though their biological activities are known, it can be evidently said that these compounds are somehow directly or indirectly responsible for the possible anti-metastatic effect on lung and colon cancer cells. However, individual testing of each identified phytochemical is required to reveal and interpret the actual involvement of these *S. repanda* compounds in anti-metastasis.

Plants in the genus *Selaginella* have diverse therapeutic potential on different cancer cells, comprising induction of apoptosis, inhibition of cell proliferation and arrest of cell cycle. There are species of *Selaginella* (*S. delicatula*, *S. tamariscina*, *S. moellendorffii*), which are known to possess potent antioxidant and antitumor activities, related to activation of apoptosis through DNA fragmentation and nucleus clotting [36], by inducing the expression of p53 and G1 arrest [37], through obstruction in fatty acid synthesis [38], inhibiting transactivation of iNOS and COX-2 via inactivating the NF- κ B and avoiding the p65 translocation [39]. Several studies revealed the potent antitumor activities by *S. uncinata* and *S. tamariscina*, while moderate activity was found against Bel-7402 and HeLa cells by *S. moellendorffii*. Furthermore, out of all *Selaginella* species in the world, *S. tamariscina* is considered one of the most biologically potent plants, which is very well-known to inhibit the growth of various cancers (breast, leukemia, gastric, lung) [36–39]. Additionally, in osteosarcoma cells, it also possesses the anti-metastatic activity by down-regulating the MMP-2 and MMP-9 secretions and increasing the TIMP-1 and TIMP-2 expressions via Akt-dependent and p38 pathways [37,38,40–42].

A similar study with *S. delicatula* revealed the cytotoxicity against cancer cells of P-388, HT-29 [43] Raji, Calu-1, lymphoma and leukemia [44], due to the presence of

phytochemicals robustaflavone and amentoflavone or its derivatives. Few other studies using *S. moellendorffii* show the growth inhibition of OVCAR-3, HeLa and FS-5 cancer cells [45,46], and anti-metastasis activity in lung cancer cells [39]. However, no studies on cytotoxicity and anti-metastasis of *S. repanda* crude extract against human cancer cells exist to date, and the mechanism of anticancer potential also remains unclear. Therefore, our study was designed to potentially identify the anti-metastatic activity of *S. repanda* crude extract against A549 and HCT-116 cancer cells with a possible molecular mechanism.

S. repanda crude extract displayed significant cytotoxic potential on both malignant A549 and HCT-116 cancer cells, and their IC_{50} were 341.1 $\mu\text{g/mL}$ and 378.8 $\mu\text{g/mL}$, respectively. We performed an in vitro wound healing assay to evaluate the effect of *S. repanda* crude extract on cell migration, as migration is an important event in the progression of metastasis and cancer. *S. repanda* crude extract significantly inhibited the migration of cancer cells in the direction of the wounded area. Such results of the present study displayed that *S. repanda* crude extract impeded cell migration, which is crucial during the early phase of wound healing. Furthermore, *S. repanda* crude extract also remarkably retarded the invasion of both malignant cancer cells. Therefore, in the present study using wound healing, adhesion and invasion assays, we have shown that *S. repanda* crude extract efficiently inhibits the metastasis in both malignant cells in vitro.

To further elucidate the possible molecular mechanism behind the anti-metastatic potential of *S. repanda* crude extract, the mRNA expression level of MMP-2 and MMP-9 genes involved in the metastasis process was investigated. Both MMP-2 and MMP-9 genes mRNA expression levels were significantly decreased in a dose-dependent manner in both malignant cells. In metastasis, MMP-9 is considered the most significant protease, and its expression is connected with the growth of local tumor, invasion and metastasis in the majority of the carcinomas [47]. Therefore, increasing evidence suggests that a particular suppression of MMP-9 activity might avert metastasis [48]. Moreover, the expression of MMP genes is mainly controlled by their activators and inhibitors at the transcriptional, post-transcriptional and at the protein level [26,49]. TIMPs are assumed to play an immense role in the inhibition of MMPs [50]. In the present study, A549 and HCT-116 cells treated with *S. repanda* crude extract collectively up-regulated the expression of TIMP-1 gene. Apart from TIMP-1, MT1-MMP is another key enzyme among the regulation of MMPs, whose overexpression has the main effect on the growth of tumors [51]. It is mainly responsible for the activation of MMP2 [52,53]. The expression of MT1-MMP gene was significantly found to be inhibited in both A549 and HCT-116 cells treated with *S. repanda* crude extract. Hence, these results revealed that anti-metastatic effect of *S. repanda* crude extract is linked to the inhibition of enzymatically degradative processes of tumor metastasis.

Natural products are produced by all organisms, but plants are the major contributors. All of these organisms co-exist in the ecosystem and interact with each other in various ways in which chemistry plays a major role. Various approaches to understand the taxonomy of plants have been evolved over the years, which include morphological, anatomical and chemotaxonomic classification. However, morphological and anatomical classification system is considered as traditional approach, whereas, the science of chemotaxonomy or chemical taxonomy is a modern approach to classify the plants, especially on the basis of their chemical constituents [54,55]. The phenolics, alkaloids, terpenoids and non-protein amino acids are the four important and widely exploited groups of compounds utilized for chemotaxonomic classification [56]. These groups of compounds exhibit a wide variation in chemical diversity, distribution and function [56,57]. From this study, *S. repanda* have been recognized as an excellent source of phytoconstituents with diverse chemical class, hence our data can also be used by ethnopharmacologists, taxonomists and ethnobotanists for chemotaxonomic importance to solve the selected taxonomical problems.

Furthermore, as we stated earlier that our study is the first study to report the phytochemistry of *S. repanda* and the identified constituents can be seen in Table 1. In the literature, around 130 chemically defined natural products are reported from 32 species of *Selaginella*, which are belonging to the classes of pigments, benzenoids, al-

kaloids, carbohydrates, coumarins, flavonoids, chromones, oxygen heterocycle, lignans, phenylpropanoids, quinoids and steroids. Here, we have discussed the phytoconstituents of other *Selaginella* species in comparison to *S. repanda*. Chao et al. 1987 identified hordenine, hordenine-[6-O-(4-hydroxy-cinnamoyl)- β -D-glucosyl]-(1,3)- α -L-rhamnoside, hordenine-O- α -L-rhamnopyranoside, -methyltyramine-O- α -L-rhamnoside as alkaloids in *S. doederleinii* [58]. Similarly, in *S. moellendorffii*, identified alkaloids were selaginelllic acid, 5-hydroxyselaginelllic acid, 5-hydroxy-N8,N8-dimethylpseudophrynaminol, N-selaginelloyl-L-phenylalanine, N-(5-hydroxyselaginelloyl)-L-phenylalanine, neoselaginelllic acid, and N-(5-Hydroxyneoselaginelloyl)-L-phenylalanine [59]. In *S. tamariscina*, adenosine and guanosine were identified as alkaloids [60]. Likewise, 4-hydroxy-benzoic acid [61], arbutin, vanillic acid, syringic acid [60,62] are identified as benzoids in *S. pulvinata* and *S. tamariscina* respectively.

Additionally, in carbohydrate class, selaginose was identified in *S. adunca*, *S. asperula*, *S. epirrhizos*, *S. galeotti*, *S. geniculata*, *S. kraussiana*, *S. marginata*, *S. parkeri*, *S. plumosa*, *S. sanguinolenta*, *S. stellate* and *S. sulcata* [63]; 2-carboxy-arabinitol was identified in *S. merten-sii* [64]; and mycose was identified in *S. pulvinata* [61]. Moreover, 8-methyl-eugenitol, uncinoside A, uncinoside B was identified as chromone in *S. uncinata* [65,66]. On the other hand, numerous flavonoids were identified in various *Selaginella* species. For example, 2,3-dihydroamentoflavone, 2'',3''-dihydroamentoflavone, tetrahydro-amentoflavone was found in *S. bryopteris* [67]; amentoflavone-7,4,7,4-tetramethylether was found in *S. moellendorffii* [68]; 4',7''-di-O-methyl-amentoflavone was found in *S. sinensis* [69]; 4',7''-di-O-methyl-amentoflavone was found in *S. willdenowii* [70]; amentoflavone was identified in various species of *Selaginella* i.e., *S. braunii*, *S. davidii*, *S. delicatula*, *S. denticulata*, *S. kraussiana*, *S. moellendorffii*, *S. pulvinata*, *S. rupestris*, *S. sanguinolenta*, *S. selaginoides*, *S. sinensis*, *S. stauntoniana*, *S. tamariscina*, *S. uncinata*, *S. willdenowii* [44,69–75]; apigenin-7-O- β -neohesperidoside, apigenin-8-C- β -D-glucopyranoside, 6,8-di-C- β -D-glucopyranosyl-apigenin, 6-C- β -D-glucopyranosyl-8-C- β -D-xylopyranosyl-apigenin, 6-C- β -D-xylopyranosyl-8-C- β -D-glucopyranosyl-apigenin was identified in *S. moellendorffii* [76,77]; 2'',3''-dihydro-4',7,7''-trimethylether-robustaflavone, 2,3-dihydro-4',7,7''-trimethylether-robustaflavone, 2'',3''-dihydro-4',7,-dimethylether-robustaflavone, 4',7-dimethylether-robustaflavone, 4'-methylether-robustaflavone was identified in *S. delicatula* [44,78]; and sumaflavone was identified in *S. tamariscina* [79].

Lignan was also found in other *Selaginella* species. 5-acethyl-dihydro-2-(3',5'-dimethoxy-4'-hydroxy-phenyl)-7-methoxybenzofuran was found in *S. tamariscina* [60]; (-)-lirioresinol A, (-)-lirioresinol B, (+)-matairesinol was found in *S. doederleinii* [18]; syringaresinol, tamariscinoside B, tamariscinoside C was found in *S. tamariscina* [60,62]. Some important pigments are also known to be found in various *Selaginella* species, i.e., selaginellin in *S. sinensis* [80], selaginellin A, selaginellin B in *S. tamariscina* [81], selaginellin C, selaginellin D, selaginellin E, selaginellin F, selaginellin G, selaginellin H in *S. pulvinata* [82–84]. Quinoids are also known to be found in some *Selaginella* species. Chrysophanic acid, emodin, physcion are identified in *S. stauntoniana* [85]; 1-methoxy-3-methylanthraquinone was identified in *S. tamariscina* [85]. Similarly, steroids are also known to be present in numerous *Selaginella* species, i.e., cholesterol, 22-dehydrocampesterol in *S. delicatula*, *S. doederleinii* [86], 3 β -16 α -dihydroxy-(5 α)-cholestan-21-oic acid in *S. pulvinata* [87], 24 α -ethyl-cholest-5-en-3 β -ol, 24 α -methyl-cholest-5-en-3 β -ol, 24 β -methyl-cholest-5-en-3 β -ol, 24 α -ethyl-cholesta-5,22-dien-3 β -ol in *S. delicatula* and *S. doederleinii* [86] and β -sitostero in *S. doederleinii*, *S. moellendorffii* and *S. pulvinata* [61,88,89]. Therefore, our results can also be correlated and compared with the possibility to identify other *Selaginella* species for their anti-metastatic and anticancer efficacy against various carcinomas, due to the presence of diverse and potential phytoconstituents.

4. Materials and Methods

4.1. Plant Material Collection and Extraction

Whole plant of *S. repanda* was collected from the wild regions of Gujarat state, India during the July–August period of 2019. The voucher specimen (BVBRC035) was deposited at Bapalal Vaidya Botanical Garden, Department of Biosciences, Veer Narmad South Gujarat University, Surat, Gujarat, India. The collected plant material was dried in an oven and then grounded into fine powder followed by storage in airtight containers. A total of 20 g of *S. repanda* powder was soaked in 85% ethanol for 24 h at 37 °C with vigorous shaking. The ethanol phase was filtered with Whatman no. 1 filter paper and then concentrated using a rotary evaporator to get the dried residue. All the assays were performed using a stock solution of crude ethanol extract.

4.2. HR-LC–MS Analysis

Phytochemistry of *S. repanda* crude extract was analyzed using UHPLC-PDA-Detector Mass Spectrophotometer (HR-L = CMS 1290 Infinity UHPLC System), Agilent Technologies®, Santa Clara, CA, USA. The liquid chromatographic system consisted of the HiP sampler, binary gradient solvent pump, column compartment and quadrupole time of flight mass spectrometer (MS Q-TOF) with the dual Agilent Jet Stream Electrospray (AJS ES) ion source. Of the sample, 10 µL was injected into the system, followed by separation in the SB-C18 column (2.1 mm × 50 mm, 1.8 µm particle size). Solvent A (1% formic acid in deionized water) and solvent B (acetonitrile) were used as solvents. A flow rate of 0.350 mL/min was used, while MS detection was performed in MS Q-TOF. Compounds were identified via their mass spectra and their unique mass fragmentation patterns. Compound Discoverer 2.1, ChemSpider and PubChem were used as the main tools for the identification of the phytochemical constituents [9].

4.3. Cell culture and Treatment

Human lung (A549) and colon (HCT-116) cancer cell lines were obtained from National Centre for Cell Science (NCCS), India and propagated in a humidified atmosphere with 5% CO₂ at 37 °C. Cells were maintained in 25 cm² flask having Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% Fetal Bovine Serum (FBS). They were grown up to 80% confluence for future analyses. Cells were treated with different concentrations of *S. repanda* crude extract (100–500 µg/mL).

4.4. Cell Viability Analysis Using MTT Assay

To determine the cytotoxicity of *S. repanda* crude extract, an MTT colorimetric assay was performed. For seeding, 96-well plates were used for both human cancer cell lines with incubation in humidified atmosphere comprising of 5% CO₂ at 37 °C up to adherence. Different concentrations of *S. repanda* crude extract (100–500 µg/mL) were then used to treat the cells for 24 h, followed by washing with PBS solution. Cells were then subjected with 100 µL of MTT solution (3-(4,5-dimethylthiazolyl-2)-2,5-diphenyltetrazolium bromide) (5 mg/mL), followed by 4-h incubation. The medium was then removed and to solubilize the formazan crystals, 100 µL of dimethyl sulfoxide (DMSO) was added. Using ELISA reader, the amount of formazan crystal was determined by measuring the absorbance at 570 nm. All assays were carried out in triplicate, and 50% cytotoxic concentration (IC₅₀) was calculated. 5-Fluorouracil was used as a positive control.

4.5. Wound Closure Assay

The effect of *S. repanda* crude extract on the motility of A-549 and HCT-116 cells was carried out using the wound healing assay. Confluent monolayer cell culture grown in 6 well plates was used to perform the assay. Cells were seeded at 1 × 10⁶ cells/mL in 3 mL final volume of growth medium. With the help of a sterile 1 mL pipette tip, an injury line was made in the central area of culture. After washing with phosphate-buffered saline (PBS), DMEM medium containing 2% FBS with and without *S. repanda* crude extract

(100–500 µg/mL) was added into the well, floating cells were discarded. Plates were then incubated for 48 h at 37 °C. Wound closure through cell migration was measured at 0, 24 and 48 h under a microscope. The cells migration towards the wound scratches was expressed as migrated cells percentage and calculated as [90]:

$$\text{Migrated cells percentage} = [(A_t = 0 \text{ h} - A_t = \Delta h) / A_t = 0 \text{ h}] \times 100 \quad (1)$$

where,

$A_t = 0 \text{ h}$ is the area of wound measured immediately after scratching

$A_t = \Delta h$ is the area of wound measured 24 or 48 h after scratching

4.6. Invasion Assay

The effect of *S. repanda* crude extract on the invasion ability of A549 and HCT-116 cells were observed using Transwell® chambers with 6.5 mm polycarbonate filters of 8 µm pore size (TCP152, Himedia®, India). A 50 µL aliquot of the Matrigel (E1270, Merck®, India) was pipetted into the upper chamber (culture insert), which was placed in a lower chamber with sterile forceps and incubated for the solidification of the gel at 37 °C for 30 min. Cells were seeded into the upper chamber with 1×10^6 cells/mL consisting of 100 µL of serum free media. Immediately this chamber was then transferred into the lower chamber consisting of 600 µL of medium (2% FBS) with and without *S. repanda* crude extract (100–500 µg/mL). The plates were then incubated for 24 h at 37 °C. Afterward, the upper chamber was taken out, and using the cotton buds, non-migrated cells on its inner surface were wiped off. Furthermore, the migrated cells on the outer surface of the upper chamber were fixed using 70% methanol for 10 min. It was then followed by staining with 0.2% crystal violet, washing with distilled water and air drying at room temperature [91]. Under an inverted microscope, invaded cells were then observed. The cells invasion through the permeable matrix gel was expressed as percentage of invading cells and calculated as:

$$\% \text{ invading cells} = [\text{Mean number of cells invading through the permeable matrix gel} / \text{Mean number of cells migrating through uncoated culture insert}] \times 100 \quad (2)$$

Mean number of cells migrating through uncoated culture insert.

4.7. Cell Adhesion Assay

The effect of *S. repanda* crude extract on the adhesion ability of A549 and HCT-116 cells was carried out according to the method described previously by Burg-Roderfeld et al., (2007) [92]. Firstly, 0.1% gelatin was coated on the surface of the 6 well culture plates and left to air dry at 37 °C for 30 min. Both A549 and HCT-116 cells (1×10^6 cells/mL) were harvested by trypsin and resuspended in a medium with and without *S. repanda* crude extract (100–500 µg/mL). Then, cells were dispensed into the respective wells and further incubated at 37 °C for 2, 4 and 6 h in 5% CO₂. At the end of incubation, attached cells were gently washed with PBS twice and counted time dependently under an inverted microscope.

4.8. Expression Levels Determination of Metastasis Related Genes

Using the TriPure Isolation Reagent (Sigma-Aldrich®, India) and according to the manufacturer's instructions, cellular RNA was isolated. RNA was quantified electrophoretically using 1.2% agarose gel, staining with ethidium bromide and visualizing under UV light. Firstly, RT-first strand synthesis kit (Qiagen®, CA, USA) was used to reverse transcribe the 1 µg of isolated RNA, and then the relative expression of metastasis related genes was determined by SYBR green based qRT PCR method (Applied Biosystems® 7500 Fast Real-Time PCR machine, CA, USA). $\Delta\Delta C_t$ method was then followed to analyze the data, and values were expressed in terms of fold change relative to control [93,94]. Cycling conditions for relative expression of genes were as follows: initial reverse transcription at 55 °C for 45 min, 1 cycle denaturation of 95 °C with 10 min hold, followed by 40 cycles of 95 °C with 15 s hold, annealing temperature at 60 °C (MMP-2, MMP-9, MT1-MMP, TIMP-1 and GAPDH) with a 60 s hold. Four pairs of primers were separately used (Table 2).

Samples were run in triplicate and their relative expression was determined by normalizing the expression of each target GAPDH.

Table 2. List of primers used for metastasis related genes [95].

Sl. No	Primer	Sequence
1	MMP-2	sense-5'-GGCCCTGTCACCTCCTGAGAT-3' antisense-5'-GGCATCCAGGTTATCGGGGA-3'
2	MMP-9	sense-5'- CGGAGCACGGAGACGGGTAT-3' antisense-5'- TGAAGGGGAAGACGCACAGC-3'
3	MT1-MMP	sense-5'-TGGGTAGCGATGAAGTCTTC-3' antisense-5'-AGTAAAGCAGTCGCTTGGGT-3'
4	TIMP-1	sense-5'- GATCCAGCGCCCAGAGAGACACC-3' antisense-5'-TTCCACTCCGGGCAGCATT-3'
5	GAPDH	sense-5'- CGAGATCCCTCCAAAATCAA-3' antisense-5'-AGGTCCACCACTGACACGTT-3'

4.9. Statistical Analysis

All the results are expressed as mean \pm SD of the number of experiments performed. A significance test was carried out among the treatments by one way ANOVA followed by Tukey's post hoc test at $p < 0.05$. Statistical analysis was conducted with software GraphPad Prism 5.0.

5. Conclusions

As per our knowledge, this study is the first one that demonstrates the anti-metastatic effect of *S. repanda* crude extract on lung and colon cancer cells. In conclusion, this study revealed that *S. repanda* crude extract exerts an inhibitory effect on various crucial steps of metastasis such as cell adhesion, invasion and migration via modulating the activities of metastasis related proteases and their activators and inhibitors. This reveals that *S. repanda* can be used/recommended as a potential anti-metastatic agent for drug development and therapy in cancer treatment.

Author Contributions: Conceptualization, M.A., M.A.K. and M.P.; methodology, M.S. (Mejdi Snoussi), A.J., A.J.S. and S.A.A.; validation, M.S. (Manojkumar Sachidanandan), M.S. (Mejdi Snoussi), W.S.H. and A.M.A.; formal analysis, M.P., M.A., M.S. (Mejdi Snoussi) and A.J.S.; investigation, R.B., A.J., S.A.A. and A.M.A.; data curation, M.P., A.J., W.S.H., S.A.A. and M.S. (Manojkumar Sachidanandan); writing—original draft preparation, M.P. and M.A.; writing—review and editing, M.A.K., M.A. and S.H.; visualization, M.P., R.B. and W.S.H.; supervision, M.A., S.H. and M.A.K.; project administration, S.H., M.A.K. and M.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research has been funded by Scientific Research Deanship at University of Ha'il Saudi Arabia through project number RG-191194.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data generated or analyzed during this study are included in this article.

Acknowledgments: I would like to thank and appreciate all the support and technical assistance provided by Mushtaq's lab at UAEU during this study. I further extend my gratitude to UoH for graciously providing numerous contributions and making all endeavors possible.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- Sung, H.; Ferlay, J.; Siegel, R.L.; Laversanne, M.; Soerjomataram, I.; Jemal, A.; Bray, F. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **2021**. [\[CrossRef\]](#)
- Coghlin, C.; I Murray, G. Current and emerging concepts in tumour metastasis. *J. Pathol.* **2010**, *222*, 1–15. [\[CrossRef\]](#)
- Adnan, M.; Khan, S.; Al-Shammari, E.; Patel, M.; Saeed, M.; Hadi, S.; Information, P.E.K.F.C. In pursuit of cancer metastasis therapy by bacteria and its biofilms: History or future. *Med. Hypotheses* **2017**, *100*, 78–81. [\[CrossRef\]](#) [\[PubMed\]](#)
- Sak, K. Chemotherapy and Dietary Phytochemical Agents. *Chemother. Res. Pr.* **2012**, *2012*, 282570. [\[CrossRef\]](#)
- Baskar, R.; Dai, J.; Wenlong, N.; Yeo, R.; Yeoh, K.-W. Biological response of cancer cells to radiation treatment. *Front. Mol. Biosci.* **2014**, *1*, 24. [\[CrossRef\]](#)
- Chikara, S.; Nagaprashantha, L.D.; Singhal, J.; Horne, D.; Awasthi, S.; Singhal, S.S. Oxidative stress and dietary phyto-chemicals: Role in cancer chemoprevention and treatment. *Cancer Lett.* **2018**, *413*, 122–134. [\[CrossRef\]](#) [\[PubMed\]](#)
- Singh, S.; Sharma, B.; Kanwar, S.S.; Kumar, A. Lead Phytochemicals for Anticancer Drug Development. *Front. Plant Sci.* **2016**, *7*, 1667. [\[CrossRef\]](#)
- Adnan, M.; Siddiqui, A.J.; Jamal, A.; Hamadou, W.S.; Awadelkareem, A.M.; Sachidanandan, M.; Patel, M. Evidence-Based Medicinal Potential and Possible Role of Selaginella in the Prevention of Modern Chronic Diseases: Ethnopharmacological and Ethnobotanical Perspective. *Rec. Nat. Prod.* **2021**, *15*, 330–355. [\[CrossRef\]](#)
- Adnan, M.; Siddiqui, A.J.; Hamadou, W.S.; Patel, M.; Ashraf, S.A.; Jamal, A.; Awadelkareem, A.M.; Sachidanandan, M.; Snoussi, M.; De Feo, V. Phytochemistry, Bioactivities, Pharmacokinetics and Toxicity Prediction of Selaginella repanda with Its Anticancer Potential against Human Lung, Breast and Colorectal Carcinoma Cell Lines. *Molecules* **2021**, *26*, 768. [\[CrossRef\]](#) [\[PubMed\]](#)
- Han, B.-H.; Chi, H.-J.; Han, Y.-N.; Ryu, K.-S. Screening on the anti-inflammatory activity of crude drugs. *Korean J. Pharmacogn.* **1972**, *3*, 205–209.
- Itokawa, H.; Mihashi, S.; Watanabe, K.; Natsumoto, H.; Hamanaka, T. Studies on the constituents of crude drugs having inhibitory activity against contraction of the ileum caused by histamine or barium chloride (1) screening test for the activity of commercially available crude drugs and the related plant materials. *Shoyakugaku Zasshi* **1983**, *37*, 223–228.
- Macfoy, C.A.; Sama, A.M. Medicinal plants in puehun district of sierra leone. *J. Ethnopharmacol.* **1983**, *8*, 215–223. [\[CrossRef\]](#)
- Han, D.S.L.; Lee, H.K. Ethnobotanical survey in Korea. In Proceedings of the Fifth Asian Symposium on Medicinal Plants and Spices, Seoul, Korea, 20–24 August 1984; Volume 5, p. 125.
- Winkelman, M. Frequently used medicinal plants in Baja California Norte. *J. Ethnopharmacol.* **1986**, *18*, 109–131. [\[CrossRef\]](#)
- Darias, V.; Bravo, L.; Rabanal, R.; Mateo, C.; Luis, R.; Pérez, A. New contribution to the ethnopharmacological study of the canary islands. *J. Ethnopharmacol.* **1989**, *25*, 77–92. [\[CrossRef\]](#)
- Ono, K.; Nakane, H.; MENG, Z.-M.; OSE, Y.; SAKAI, Y.; MIZUNO, M. Differential inhibitory effects of various herb ex-tracts on the activities of reverse transcriptase and various deoxyribonucleic acid (DNA) polymerases. *Chem. Pharm. Bull.* **1989**, *37*, 1810–1812. [\[CrossRef\]](#) [\[PubMed\]](#)
- Meng, Z.; Sakai, Y.; Ose, Y.; Sato, T.; Nagase, H.; Kito, H.; Matsuda, H.; Sato, M.; Mizuno, M. Antimutagenic activity of medical plants in traditional Chinese medicines. *Mutat. Res. Mutagen. Relat. Subj.* **1988**, *203*, 378–379. [\[CrossRef\]](#)
- Lin, R.; Skaltsounis, A.-L.; Seguin, E.; Tillequin, F.; Koch, M. Phenolic Constituents of Selaginella doederleinii. *Planta Medica* **1994**, *60*, 168–170. [\[CrossRef\]](#)
- De Sá, P.G.S.; Nunes, X.P.; De Lima, J.T.; Filho, J.A.D.S.; Fontana, A.P.; Siqueira, J.D.S.; Quintans-Júnior, L.J.; Damasceno, P.K.F.; Branco, C.R.C.; Branco, A.; et al. Antinociceptive effect of ethanolic extract of Selaginella convoluta in mice. *BMC Complement. Altern. Med.* **2012**, *12*, 187. [\[CrossRef\]](#)
- Almeida, J.R.G.; de Sa, P.G.S.; Macedo, L.A.R.; Filho, J.A.; Oliviera, V.R.; Filho, J.M.B. Phytochemistry of the genus Selaginella (Selaginellaceae). *J. Med. Plants Res.* **2013**, *7*, 1858–1868.
- Harbone, J.B.; Williams, C.A. ChemInform Abstract: Advances in Flavonoid Research since 1992. *Phytochemistry* **2010**, *32*, 481–504. [\[CrossRef\]](#)
- Pellati, F.; Benvenuti, S.; Magro, L.; Melegari, M.; Soragni, F. Analysis of phenolic compounds and radical scavenging activity of Echinacea spp. *J. Pharm. Biomed. Anal.* **2004**, *35*, 289–301. [\[CrossRef\]](#)
- Lan, L.; Wang, Y.; Pan, Z.; Wang, B.; Yue, Z.; Jiang, Z.; Li, L.; Wang, C.; Tang, H. Rhamnetin induces apoptosis in human breast cancer cells via the miR-34a/Notch-1 signaling pathway. *Oncol. Lett.* **2018**, *17*, 676–682. [\[CrossRef\]](#) [\[PubMed\]](#)
- Lee, Y.-C.; Lin, H.-H.; Hsu, C.-H.; Wang, C.-J.; Chiang, T.-A.; Chen, J.-H. Inhibitory effects of andrographolide on migration and invasion in human non-small cell lung cancer A549 cells via down-regulation of PI3K/Akt signaling pathway. *Eur. J. Pharmacol.* **2010**, *632*, 23–32. [\[CrossRef\]](#) [\[PubMed\]](#)
- Yeh, C.-B.; Hsieh, M.-J.; Lin, C.-W.; Chiou, H.-L.; Lin, P.-Y.; Chen, T.-Y.; Yang, S.-F. The antimetastatic effects of resveratrol on hepatocellular carcinoma through the downregulation of a metastasis-associated protease by SP-1 modulation. *PLoS ONE* **2013**, *8*, e56661. [\[CrossRef\]](#)

26. Yang, S.-F.; Chen, M.-K.; Hsieh, Y.-S.; Yang, J.-S.; Zavras, A.-I.; Hsieh, Y.-H.; Su, S.-C.; Kao, T.-Y.; Chen, P.-N.; Chu, S.-C. Antimetastatic effects of *Terminalia catappa* L. on oral cancer via a down-regulation of metastasis-associated proteases. *Food Chem. Toxicol.* **2010**, *48*, 1052–1058. [\[CrossRef\]](#)
27. Giannelli, G.; Bergamini, C.; Fransvea, E.; Marinosci, F.; Quaranta, V.; Antonaci, S. Human Hepatocellular Carcinoma (HCC) Cells Require Both $\alpha 3\beta 1$ Integrin and Matrix Metalloproteinases Activity for Migration and Invasion. *Lab. Investig.* **2001**, *81*, 613–627. [\[CrossRef\]](#)
28. Patel, M.; Sachidanandan, M.; Adnan, M. Serine arginine protein kinase 1 (SRPK1): A moonlighting protein with theranostic ability in cancer prevention. *Mol. Biol. Rep.* **2018**, *46*, 1487–1497. [\[CrossRef\]](#)
29. Liotta, L.A. An attractive force in metastasis. *Nat. Cell Biol.* **2001**, *410*, 24–25. [\[CrossRef\]](#)
30. Reddy, M.N.; Adnan, M.; Alreshidi, M.M.; Saeed, M.; Patel, M. Evaluation of Anticancer, Antibacterial and Antioxidant Properties of a Medicinally Treasured Fern *Tectaria coadunata* with its Phytoconstituents Analysis by HR-LCMS. *Anti-Cancer Agents Med. Chem.* **2020**, *20*, 1845–1856. [\[CrossRef\]](#)
31. Siddiqui, A.J.; Danciu, C.; Ashraf, S.A.; Moin, A.; Singh, R.; Alreshidi, M.; Patel, M.; Jahan, S.; Kumar, S.; Alkhinjar, M.I. Plants-derived biomolecules as potent antiviral phytomedicines: New insights on ethnobotanical evidences against coronavirus. *Plants* **2020**, *9*, 1244. [\[CrossRef\]](#)
32. Patel, M.; Ashraf, M.S.; Siddiqui, A.J.; Ashraf, S.A.; Sachidanandan, M.; Snoussi, M.; Adnan, M.; Hadi, S. Profiling and Role of Bioactive Molecules from *Puntius sophore* (Freshwater/Brackish Fish) Skin Mucus with Its Potent Antibacterial, Antiadhesion, and Antibiofilm Activities. *Biomolecules* **2020**, *10*, 920. [\[CrossRef\]](#) [\[PubMed\]](#)
33. Surti, M.; Patel, M.; Adnan, M.; Moin, A.; Ashraf, S.A.; Siddiqui, A.J.; Snoussi, M.; Deshpande, S.; Reddy, M.N. Ili-maquione (marine sponge metabolite) as a novel inhibitor of SARS-CoV-2 key target proteins in comparison with suggested COVID-19 drugs: Designing, docking and molecular dynamics simulation study. *RSC Adv.* **2020**, *10*, 37707–37720. [\[CrossRef\]](#)
34. Adnan, M.; Patel, M.; Deshpande, S.; Alreshidi, M.; Siddiqui, A.J.; Reddy, M.N.; Emira, N.; De Feo, V. Effect of *Adiantum philippense* Extract on Biofilm Formation, Adhesion with Its Antibacterial Activities Against Foodborne Pathogens, and Characterization of Bioactive Metabolites: An in vitro-in silico Approach. *Front. Microbiol.* **2020**, *11*, 823. [\[CrossRef\]](#)
35. Mseddi, K.; Alimi, F.; Noumi, E.; Veettil, V.N.; Deshpande, S.; Adnan, M.; Hamdi, A.; Elkahoui, S.; Alghamdi, A.; Kadri, A.; et al. *Thymus musilii* Velen. as a promising source of potent bioactive compounds with its pharmacological properties: In vitro and in silico analysis. *Arab. J. Chem.* **2020**, *13*, 6782–6801. [\[CrossRef\]](#)
36. Ahn, S.-H.; Mun, Y.-J.; Lee, S.-W.; Kwak, S.; Choi, M.-K.; Baik, S.-K.; Kim, Y.-M.; Woo, W.-H. *Selaginella tamariscina* Induces Apoptosis via a Caspase-3-Mediated Mechanism in Human Promyelocytic Leukemia Cells. *J. Med. Food* **2006**, *9*, 138–144. [\[CrossRef\]](#)
37. Lee, I.-S.; Nishikawa, A.; Furukawa, F.; Kasahara, K.-I.; Kim, S.-U. Effects of *Selaginella tamariscina* on in vitro tumor cell growth, p53 expression, G1 arrest and in vivo gastric cell proliferation. *Cancer Lett.* **1999**, *144*, 93–99. [\[CrossRef\]](#)
38. Lee, J.S.; Lee, M.S.; Oh, W.K.; Sul, J.Y. Fatty acid synthase inhibition by amentoflavone induces apoptosis and antiproliferation in human breast cancer cells. *Biol. Pharm. Bull.* **2009**, *32*, 1427–1432. [\[CrossRef\]](#)
39. Woo, E.-R.; Pokharel, Y.R.; Yang, J.W.; Lee, S.Y.; Kang, K.W. Inhibition of nuclear factor- κ B activation by 2', 8''-biapigenin. *Biol. Pharm. Bull.* **2006**, *29*, 976–980. [\[CrossRef\]](#)
40. Setyawan, A.D. Review: Natural products from Genus *Selaginella* (Selaginellaceae). *Nusant. Biosci.* **2016**, *3*, 3. [\[CrossRef\]](#)
41. Yang, J.-S.; Lin, C.-W.; Hsieh, Y.-S.; Cheng, H.-L.; Lue, K.-H.; Yang, S.-F.; Lu, K.-H. *Selaginella tamariscina* (Beauv.) possesses antimetastatic effects on human osteosarcoma cells by decreasing MMP-2 and MMP-9 secretions via p38 and Akt signaling pathways. *Food Chem. Toxicol.* **2013**, *59*, 801–807. [\[CrossRef\]](#)
42. Yang, J.-S.; Lin, C.-W.; Hsin, C.-H.; Hsieh, M.-J.; Chang, Y.-C. *Selaginella tamariscina* Attenuates Metastasis via Akt Pathways in Oral Cancer Cells. *PLoS ONE* **2013**, *8*, e68035. [\[CrossRef\]](#) [\[PubMed\]](#)
43. Chen, J.-J.; Duh, C.-Y.; Chen, J.-F. New cytotoxic biflavonoids from *Selaginella delicatula*. *Planta Med.* **2005**, *71*, 659–665. [\[CrossRef\]](#) [\[PubMed\]](#)
44. Lin, L.-C.; Kuo, Y.-C.; Chou, C.-J. Cytotoxic Biflavonoids from *Selaginella delicatula*. *J. Nat. Prod.* **2000**, *63*, 627–630. [\[CrossRef\]](#)
45. Sun, C.-M.; Syu, W.-J.; Huang, Y.-T.; Chen, C.-C.; Ou, J.-C. Selective cytotoxicity of ginkgetin from *Selaginella moellendorffii*. *J. Nat. Prod.* **1997**, *60*, 382–384. [\[CrossRef\]](#) [\[PubMed\]](#)
46. Su, Y.; Sun, C.-M.; Chuang, H.-H.; Chang, P.-T. Studies on the cytotoxic mechanisms of ginkgetin in a human ovarian adenocarcinoma cell line. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **2000**, *362*, 82–90. [\[CrossRef\]](#)
47. Sato, H.; Takino, T.; Okada, Y.; Cao, J.; Shinagawa, A.; Yamamoto, E.; Seiki, M. A matrix metalloproteinase expressed on the surface of invasive tumour cells. *Nat. Cell Biol.* **1994**, *370*, 61–65. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Yeh, C.-B.; Hsieh, M.-J.; Hsieh, Y.-H.; Chien, M.-H.; Chiou, H.-L.; Yang, S.-F. Antimetastatic effects of norcantharidin on hepatocellular carcinoma by transcriptional inhibition of MMP-9 through modulation of NF- κ B activity. *PLoS ONE* **2012**, *7*, e31055. [\[CrossRef\]](#)
49. Ghiso, J.A.A.; Alonso, D.F.; Fariás, E.F.; Gomez, D.E.; Joffé, E.B.D.K. Deregulation of the signaling pathways controlling urokinase production. Its relationship with the invasive phenotype. *JBC J. Biol. Inorg. Chem.* **1999**, *263*, 295–304. [\[CrossRef\]](#)
50. Hwang, E.-S.; Lee, H.J. Allyl isothiocyanate and its N-acetylcysteine conjugate suppress metastasis via inhibition of invasion, migration, and matrix metalloproteinase-2/-9 activities in SK-Hep1 human hepatoma cells. *Exp. Biol. Med.* **2006**, *231*, 421–430. [\[CrossRef\]](#) [\[PubMed\]](#)

51. Määttä, M.; Soini, Y.; Liakka, A.; Autio-Harmainen, H. Differential expression of matrix metalloproteinase (MMP)-2, MMP-9, and membrane type 1-MMP in hepatocellular and pancreatic adenocarcinoma: Implications for tumor progression and clinical prognosis. *Clin. Cancer Res.* **2000**, *6*, 2726–2734.
52. Ogata, R.; Torimura, T.; Kin, M.; Ueno, T.; Tateishi, Y.; Kuromatsu, R.; Shimauchi, Y.; Sakamoto, M.; Tamaki, S.; Sata, M.; et al. Increased expression of membrane type 1 matrix metalloproteinase and matrix metalloproteinase-2 with tumor dedifferentiation in hepatocellular carcinomas. *Hum. Pathol.* **1999**, *30*, 443–450. [\[CrossRef\]](#)
53. Tam, E.M.; Moore, T.R.; Butler, G.S.; Overall, C.M. Characterization of the Distinct Collagen Binding, Helicase and Cleavage Mechanisms of Matrix Metalloproteinase 2 and 14 (Gelatinase A and MT1-MMP). *J. Biol. Chem.* **2004**, *279*, 43336–43344. [\[CrossRef\]](#)
54. Reynolds, T. The evolution of chemosystematics. *Phytochemistry* **2007**, *68*, 2887–2895. [\[CrossRef\]](#) [\[PubMed\]](#)
55. Larsen, T.O.; Smedsgaard, J.; Nielsen, K.F.; Hansen, M.E.; Frisvad, J.C. Phenotypic taxonomy and metabolite profiling in microbial drug discovery. *Nat. Prod. Rep.* **2005**, *22*, 672–695. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Fellows, L.; Smith, P.M. The Chemotaxonomy of Plants. *Kew Bull.* **1978**, *32*, 806. [\[CrossRef\]](#)
57. Hegnauer, R. Phytochemistry and plant taxonomy—An essay on the chemotaxonomy of higher plants. *Phytochemistry* **1986**, *25*, 1519–1535. [\[CrossRef\]](#)
58. Chao, L.R.; Seguin, E.; Tillequin, F.; Koch, M. New Alkaloid Glycosides from *Selaginella doederleinii*. *J. Nat. Prod.* **1987**, *50*, 422–426. [\[CrossRef\]](#)
59. Wang, Y.-H.; Long, C.-L.; Yang, F.-M.; Wang, X.; Sun, Q.-Y.; Wang, H.-S.; Shi, Y.-N.; Tang, G.-H. Pyrrolidinoindoline Alkaloids from *Selaginella moellendorffii*. *J. Nat. Prod.* **2009**, *72*, 1151–1154. [\[CrossRef\]](#)
60. Zheng, X.-K.; Bi, Y.-F.; Feng, W.-S.; Shi, S.-P.; Wang, J.-F.; Niu, J.-Z. Study on chemical constituents of *Selaginella tamariscina* (Beauv.) Spring. *Acta Pharm. Sin.* **2004**, *39*, 266–268.
61. Zheng, X.; Liao, D.F.; Zhu, B.Y.; Tuo, Q.H.; Xu, Y.L. Study on chemical constituents of *Selaginella pulvinata*. *Zhongcaoyao* **2001**, *32*, 17–18.
62. Bi, Y.-F.; Zheng, X.-K.; Feng, W.-S.; Shi, S.-P. Isolation and structural identification of chemical constituents from *Selaginella tamariscina* (Beauv.) Spring. *Acta Pharm. Sin.* **2004**, *39*, 41–45.
63. Fischer, M.; Kandler, M. Identifizierung von selaginose und deren verbreitung in der gattung *Selaginella*. *Phytochemistry* **1975**, *14*, 2629–2633. [\[CrossRef\]](#)
64. Moore, B.D.; Isidoro, E.; Seemann, J.R. Distribution of 2-carboxyarabinitol among plants. *Phytochemistry* **1993**, *34*, 703–707. [\[CrossRef\]](#)
65. Ma, L.-Y.; Ma, S.-C.; Wei, F.; Lin, R.-C.; But, P.P.-H.; Lee, S.H.-S.; Lee, S.F. Uncinoside A and B, Two New Antiviral Chromone Glycosides from *Selaginella uncinata*. *Chem. Pharm. Bull.* **2003**, *51*, 1264–1267. [\[CrossRef\]](#) [\[PubMed\]](#)
66. Ma, L.Y.; Wei, F.; Ma, S.C.; Lin, R.C. Two new chromone glycosides from *Selaginella uncinata*. *Chin. Chem. Lett.* **2002**, *13*, 748–751.
67. Kunert, O.; Swamy, R.C.; Kaiser, M.; Presser, A.; Buzzi, S.; Rao, A.A.; Schühly, W. Antiplasmodial and leishmanicidal activity of biflavonoids from Indian *Selaginella bryopteris*. *Phytochem. Lett.* **2008**, *1*, 171–174. [\[CrossRef\]](#)
68. Cao, Y.; Tan, N.H.; Chen, J.J.; Zeng, G.Z.; Ma, Y.B.; Wu, Y.P.; Yan, H.; Yang, J.; Lu, L.F.; Wang, Q. Bioactive flavones and biflavones from *Selaginella moellendorffii* Hieron. *Fitoterapia* **2010**, *81*, 253–258. [\[CrossRef\]](#)
69. Ma, S.-C.; But, P.P.-H.; Ooi, V.E.-C.; He, Y.-H.; Lee, S.H.-S.; Lee, S.-F.; Lin, R.-C. Antiviral Amentoflavone from *Selaginella sinensis*. *Biol. Pharm. Bull.* **2001**, *24*, 311–312. [\[CrossRef\]](#)
70. Silva, G.L.; Chai, H.; Gupta, M.P.; Farnsworth, N.R.; Cordell, G.A.; Pezzuto, J.M.; Beecher, C.W.; Kinghorn, A.D. Cytotoxic biflavonoids from *Selaginella willdenowii*. *Phytochemistry* **1995**, *40*, 129–134. [\[CrossRef\]](#)
71. López-Sáez, J.A.; Alonso, M.J.P.-; Negueruela, A.V. Biflavonoids of *Selaginella denticulata* Growing in Spain. *Z. Nat. C* **1994**, *49*, 267–270. [\[CrossRef\]](#)
72. Qasim, M.A.; Roy, S.K.; Kamil, M.; Ilyas, M. Phenolic constituents of Selaginellaceae. *Indian J. Chem.* **1985**, *24*, 220.
73. Chakravarthy, B.K.; Rao, Y.V.; Gambhir, S.S.; Gode, K.D. Isolation of Amentoflavone from *Selaginella rupestris* and its Pharmacological Activity on Central Nervous System, Smooth Muscles and Isolated Frog Heart Preparations. *Planta Med.* **1981**, *43*, 64–70. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Huneck, S.; Khaidav, T. Amentoflavone from *Selaginella sanguinolenta*. *Die Pharm.* **1985**, *40*, 431.
75. Lee, I.R.; Song, J.Y.; Lee, Y.S. Cytotoxicity of folkloric medicines in murine and human cancer cells. *Korean J. Pharmacogn.* **1992**, *23*, 132–136.
76. Feng, W.S.; Li, K.K.; Zheng, X.K. A new norlignan lignanoside from *Selaginella moellendorffii* Hieron. *Acta Pharm. Sin. B* **2011**, *1*, 36–39. [\[CrossRef\]](#)
77. Zhu, T.M.; Chen, K.L.; Zhou, W.B. A new flavones glycoside from *Selaginella moellendorffii* Hieron. *Chin. Chem. Lett.* **2008**, *19*, 1456–1458. [\[CrossRef\]](#)
78. Lin, L.C.; Chou, C.J. Three new biflavonoids from *Selaginella delicatula*. *Chin. J. Pharm.* **2000**, *52*, 211–218.
79. Yang, J.W.; Pokharel, Y.R.; Kim, M.-R.; Woo, E.-R.; Choi, H.K.; Kang, K.W. Inhibition of inducible nitric oxide synthase by sumafavone isolated from *Selaginella tamariscina*. *J. Ethnopharmacol.* **2006**, *105*, 107–113. [\[CrossRef\]](#)
80. Zhang, L.-P.; Liang, Y.-M.; Wei, X.-C.; Cheng, D.-L. A New Unusual Natural Pigment from *Selaginella sinensis* and Its Noticeable Physicochemical Properties. *J. Org. Chem.* **2007**, *38*, 3824–3921. [\[CrossRef\]](#)
81. Cheng, X.-L.; Ma, S.-C.; Yu, J.-D.; Yang, S.-Y.; Xiao, X.-Y.; Hu, J.-Y.; Lu, Y.; Shaw, P.-C.; But, P.P.-H.; Lin, R.-C. Selaginellin A and B, Two Novel Natural Pigments Isolated from *Selaginella tamariscina*. *Chem. Pharm. Bull.* **2008**, *56*, 982–984. [\[CrossRef\]](#)

82. Tan, G.-S.; Xu, K.-P.; Li, F.-S.; Wang, C.-J.; Li, T.-Y.; Hu, C.-P.; Shen, J.; Zhou, Y.-J.; Li, Y.-J. Selaginellin C, a new natural pigment from *Selaginella pulvinata* Maxim (Hook et Grev.). *J. Asian Nat. Prod. Res.* **2009**, *11*, 1001–1004. [[CrossRef](#)] [[PubMed](#)]
83. Cao, Y.; Chen, J.J.; Tan, N.H.; Oberer, L.; Wagner, T.; Wu, Y.P.; Zeng, G.Z.; Yan, H.; Wang, Q. Antimicrobial selaginellin derivatives from *Selaginella pulvinata*. *Bioorganic Med. Chem. Lett.* **2010**, *20*, 2456–2460. [[CrossRef](#)] [[PubMed](#)]
84. Cao, Y.; Yang, J.; Wang, Q.; Chen, J.-J.; Tan, N.-H.; Wu, Y.-P. Structure determination of selaginellins G and H from *Selaginella pulvinata* by NMR spectroscopy. *Magn. Reson. Chem.* **2010**, *48*, 656–659. [[CrossRef](#)] [[PubMed](#)]
85. Liu, J.-F.; Xu, K.-P.; Li, F.-S.; Shen, J.; Hu, C.-P.; Zou, H.; Yang, F.; Liu, G.-R.; Xiang, H.-L.; Zhou, Y.-J.; et al. A New Flavonoid from *Selaginella tamariscina* (Beauv.) Spring. *Chem. Pharm. Bull.* **2010**, *58*, 549–551. [[CrossRef](#)]
86. Chiu, P.-L.; Patterson, G.W.; Salt, T.A. Sterol composition of pteridophytes. *Phytochemistry* **1988**, *27*, 819–822. [[CrossRef](#)]
87. Zheng, X.; Du, J.; Xu, Y.; Zhu, B.; Liao, D. A new steroid from *Selaginella pulvinata*. *Fitoterapia* **2007**, *78*, 598–599. [[CrossRef](#)] [[PubMed](#)]
88. Chen, P.; Sun, J.Y.; Xie, N.G.; Shi, Y.G. Chemical constituents of daeycai (*Selaginella doederleinii*). *Zhongcaoyao* **1995**, *26*, 397–399.
89. Che, D.H.; Yu, J.G. Analysis on the chemical constituents of jiangnanjuanbai (*Selaginella moellendorffii* Hieron). *Chung Tsao Yao* **1986**, *17*, 4.
90. Baraya, Y.S.; Wong, K.K.; Yaacob, N.S. *Strobilanthes crispus* inhibits migration, invasion and metastasis in breast cancer. *J. Ethnopharmacol.* **2019**, *233*, 13–21. [[CrossRef](#)] [[PubMed](#)]
91. Justus, C.R.; Leffler, N.; Ruiz-Echevarria, M.; Yang, L.V. In vitro Cell Migration and Invasion Assays. *J. Vis. Exp.* **2014**, e51046. [[CrossRef](#)]
92. Burg-Roderfeld, M.; Roderfeld, M.; Wagner, S.; Henkel, C.; Grötzinger, J.; Roeb, E. MMP-9-hemopexin domain hampers adhesion and migration of colorectal cancer cells. *Int. J. Oncol.* **2007**, *30*, 985–992. [[CrossRef](#)] [[PubMed](#)]
93. Adnan, M.; Morton, G.; Hadi, S. Analysis of rpoS and bolA gene expression under various stress-induced environments in planktonic and biofilm phase using $2^{-\Delta\Delta CT}$ method. *Mol. Cell. Biochem.* **2011**, *357*, 275–282. [[CrossRef](#)] [[PubMed](#)]
94. Siddiqui, A.J.; Bhardwaj, J.; Goyal, M.; Prakash, K.; Adnan, M.; Alreshidi, M.M.; Patel, M.; Soni, A.; Redman, W. Immune responses in liver and spleen against *Plasmodium yoelii* pre-erythrocytic stages in Swiss mice model. *J. Adv. Res.* **2020**, *24*, 29–41. [[CrossRef](#)] [[PubMed](#)]
95. Byambaragchaa, M.; De La Cruz, J.; Yang, S.H.; Hwang, S.-G. Anti-metastatic Potential of Ethanol Extract of *Saussurea involucreata* against Hepatic Cancer in vitro. *Asian Pac. J. Cancer Prev.* **2013**, *14*, 5397–5402. [[CrossRef](#)] [[PubMed](#)]